

1 **Can the effects of anthropogenic pressures and environmental variability**
2 **on nekton fauna be detected in fishery data? Insights from the monitoring**
3 **of the artisanal fishery within the Venice lagoon**
4

5 Zucchetta M^{1*}, Scapin L¹, Cavararo F¹, Pranovi F¹, Franco A², and Franzoi P¹

6 1 CEMAS – Centre for Estuarine and coastal Marine Sciences. University Ca' Foscari Venice

7 2 Institute of Estuarine & Coastal Studies. University of Hull

8 <https://doi.org/10.1007/s12237-015-0064-y>

9

10

11 * Corresponding Author: Matteo Zucchetta, Department of Environmental Sciences, Informatics and Statistics,
12 University Ca' Foscari Venice. Castello 2737/b, 30122 Venice, Italy.

13 e-mail: matzuc@unive.it

14

1. Introduction

Transitional waters are highly productive ecosystems that are affected by both a naturally high variability in environmental conditions and multiple anthropogenic pressures (Costanza et al., 1997; Elliott and Quintino, 2007; Vasconcelos et al., 2007). In order to effectively manage driving forces on European transitional water ecosystems, and thus preserve the services provided, the Water Framework Directive (WFD; Directive 2000/60/EC) requires the evaluation of ecological status, based on indices sensitive to anthropogenic pressures rather than to natural variability (Uriarte and Borja, 2009).

Fish are an ecologically and economically important component of transitional waters potentially influenced by the alteration of environmental conditions (Elliott and Dewailly, 1995; Elliott et al., 2007). As a result, several fish-based multi-metric indices have been developed to assess the ecological status of these ecosystems in Europe, and more recently anthropogenic pressures have been additionally incorporated in order to validate such indices (Aubry and Elliott, 2006; Cabral et al., 2012; Fonseca et al., 2013; Pasquaud et al., 2013). This process allows the evaluation of the sensitivity of fish-based multi-metric indices to human pressures and can promote the development of ecological status assessment methods that are fully compliant with WFD requirements (Hering et al., 2010). In this light, the choice of metrics to be included in an index is particularly relevant, as their sensitivity to human alteration should be verified (i.e. the metric should show a decrease with increasing pressure values), taking into account natural variability of the system while keeping such a relationship ecologically interpretable (Cabral et al., 2012; Pérez-Domínguez et al., 2012; Schoolmaster et al., 2012, 2013a, 2013b). In transitional waters, several human activities can generate pressures which may in turn lead to a number of direct or indirect disturbances to fish populations. The main anthropogenic pressures are related with chemical pollution, physical changes, energy and thermal pollution, radioactivity and biological pollution (species invasions and pathogens) (Marchand et al., 2002). As Vasconcelos et al. (2007) exemplified, multiple anthropogenic activities can lead to degradation of water and sediment quality (e.g. wastewater discharges and use of chemicals in agriculture), habitat loss (e.g. bank reclamation) or reduction of prey availability (e.g. sediment management). All these can potentially affect nekton fauna at various levels of biological organisation, e.g. by directly increasing mortality of individuals or indirectly causing shifts in community composition through alterations of ecosystem processes.

42 Fishing is one of the human activities which, while taking advantage of the high productivity of transitional
43 water ecosystems, can also have a role in generating system pressures for example by removing nektonic
44 biomass or by leading to modifications to the habitat (McHugh, 1967; Nixon, 1982; Blaber et al., 2000; Elliott,
45 2002). On the other hand, the species composition and biomass of fishery landings strongly depend on the
46 ecological status of the system (Pérez-Ruzafa and Marcos, 2012). Although Mediterranean coastal lagoons
47 support important fishery activities and maintain aquaculture exploitation (Kapetsky and Lasserre, 1984;
48 Ardizzone et al., 1988), the role of fishery as a cause of pressure for fish communities is poorly investigated
49 in such ecosystems. In Mediterranean transitional waters, fixed gears such as fyke nets and fishing barriers at
50 the sea inlets are probably the most important fishery techniques (Cataudella and Ferlin, 1984; Ardizzone et
51 al., 1988; Chauvet, 1988; Pérez-Ruzafa and Marcos, 2012). Such fisheries take advantage of fish movement
52 within or between the transitional system and the adjacent marine and freshwater areas, and are highly seasonal
53 due to the presence and migration of different species (Granzotto et al., 2001; Provincia di Venezia, 2009).

54 Monitoring of artisanal fisheries can provide useful information on fish assemblages or on the status and
55 evolution of some populations, provided sampling effects are accounted for (e.g gear characteristics and
56 selectivity) (Malavasi et al., 2004a; Provincia di Venezia, 2009; Pranovi et al., 2013). Nekton data from fishery
57 monitoring could be in some cases routinely collected, enhancing the possibility of gathering a large and robust
58 dataset. Moreover, if fishery data could be also used to contribute to the assessment of ecological status, it
59 would help in containing sampling costs related to scientific monitoring.

60 In this study, data were analysed from a pluriennial monitoring plan of the artisanal fishery (fyke nets) in the
61 Venice lagoon in order to test the effects of human pressures on nekton assemblage, namely fish and
62 invertebrates. We followed a model-based approach (Warton et al., 2014) in order to test *a priori*-formulated
63 hypotheses about the different role of natural variability, anthropogenic pressures and the artisanal fishery in
64 affecting the lagoon nekton assemblage, and to understand whether monitoring of the artisanal fishery can be
65 used to assess the relationship between nekton assemblage and anthropogenic pressures in transitional waters.
66 The issues addressed in this study represent a crucial step for setting up an ecological status evaluation system
67 based on monitoring of the local fishery. This could be important to optimise the effort of collecting field data
68 and to harmonise different monitoring programmes.

69 **2. Materials and methods**

70 **2.1. Venice lagoon and study areas**

71 The Venice lagoon is a large coastal lagoon (about 550 km²) in the North-Adriatic Sea (Italy) (Figure 1) It is
72 a microtidal system with mean tide amplitude of 1m (Umgiesser et al., 2004). It is a shallow system comprised
73 of three sub-basins delimited by two main watersheds (Solidoro et al., 2004). There is a high spatial and
74 temporal variability in morphological and physico-chemical parameters, as well as a mosaic of habitats,
75 including saltmarshes, seagrass meadows, bare or sparsely vegetated mudflats and sandflats (Franzoi et al.,
76 2010; Solidoro et al., 2010).

77 For the purpose of this study, five broad homogeneous areas were identified in the lagoon, based on the degree
78 of confinement (Solidoro et al., 2004), main habitat types, salinity and sediment characteristics: Chioggia (CH),
79 Ca' Zane (CZ), Lido (LD), Lago dei Teneri (LT) and Ponte della Libertà (PL) (Figure 1). CZ and PL both
80 experience high water residence times (>20 days) with euhaline conditions and predominance of muddy
81 substrata often covered by macroalgae, with some marsh areas only within CZ. LT is also very confined (>20
82 days water residence time), but it shows polyhaline conditions and dominance of saltmarsh habitats, with
83 presence of some bare or macroalgae-covered mudflats. LD and CH are characterised by euhaline conditions,
84 low residence times (<7 days) and seagrass-dominated sandy habitats.

85

86 **Figure 1 about here.**

87

88 **2.2. Monitoring programme and nekton assemblage**

89 Fyke nets represent the most important gear type of the fishery of the Venice lagoon (Granzotto et al., 2004;
90 Provincia di Venezia, 2009; Pranovi et al., 2013). Fyke nets employed here consist of a barrier of about 50 m
91 in length and 1.3 m in height, with a mesh size of 0.6 cm, which guides the fish towards four cone shaped,
92 unbaited traps (Malavasi et al., 2004a). Artisanal fishery activity in the five lagoon areas was monitored on a
93 monthly basis in two seasons (spring and autumn), corresponding to the two major fishing periods (Provincia
94 di Venezia, 2009). Data were collected during the years 2001 to 2003 and 2009 to 2013 in order to obtain a
95 representative view of species composition and biomass of nekton assemblage inside the lagoon. For each
96 study area an average number of 79.38 (Standard Deviation: 42.25) traps were inspected during each season.

97 The content of the traps was inspected on board and samples were collected to confirm identification in the
98 laboratory. Catches (including target species and by catch) were identified at species level and weighted (± 1
99 g), in order to obtain the cumulative biomass per species. This was expressed in terms of catch per unit effort
100 (CPUE), i.e. catch (in g) was standardised per trap and temporal unit, the latter accounting for the number of
101 days since the previous visit by the fishermen (Pranovi et al., 2013). Species richness (S) and biomass (B) were
102 calculated for each sample.

103 Both fish and cephalopods were considered due to their relevance to the local fisheries (Granzotto et al., 2004;
104 Provincia di Venezia, 2009; Pranovi et al., 2013). The green crab, *Carcinus aestuarii*, although representing a
105 very important target of the fishery (Pranovi et al., 2013), was not considered in the analyses, due to its benthic
106 habits and to peculiar fishing practices, based on the selection of the moulting specimens (Matozzo et al., 2013;
107 Pranovi et al., 2013). While decapods (including *Crangon crangon*, *Melicertus kerathurus*, *Processa*
108 *macrophthalma* and a number of species belonging to the genus *Palaemon*) can be relevant to the local fishery,
109 these were only recorded from 2009 onwards, and therefore they were excluded from the analysis (a
110 preliminary exploration of the 2009-2013 data indicated their minor contribution to the results).

111 The recorded species were categorised according to the role played with respect to the artisanal fishery: target
112 species, directly pursued by the fishery; incidental species, i.e. non-target species incidentally caught and with
113 commercial value; discarded species, with no commercial value (Pranovi et al., 2013). In addition, Estuarine
114 Use Functional Groups (EUFG) defining the main ecological utilisation of the lagoon by the species were
115 adopted and modified after Potter et al. (2013). The attribution of EUFG to the species was undertaken taking
116 into account the specific use of the Venice lagoon (Malavasi et al., 2004a; Franco et al., 2008; Franzoi et al.,
117 2010), as well as the type of habitat used by the species (e.g. some species were classified as marine stragglers,
118 and not estuarine residents or estuarine opportunists, due to their exclusive use of marine-like habitats included
119 within the lagoon boundaries, even if they were found frequently and with relatively high abundances). Feeding
120 Mode Functional Groups (FMFG) were adapted from Franco et al. (2008, 2009b), in order to characterise the
121 species in terms of feeding behaviour. The attribution of FMFG to the species was carried out taking into
122 account the life stage predominantly found in fyke nets. As one species can be allocated to multiple FMFG,
123 the species contribution to a single guild was assigned in proportion to the relevance of each feeding mode for
124 the species, by identifying the importance (%) of different guild allocations within the diet, on the basis of the

125 literature (“www.fishbase.org”; Froese and Pauly, 2015) and of available data for the Venice lagoon (Franzoi,
126 unpublished data). Guilds for nektonic invertebrates were also defined on the basis of available information in
127 the scientific literature (Blanc and Daguzan, 1998; Pinczon Du Sel et al., 2000; Alves et al., 2006; Palomares
128 and Pauly, 2014). A complete list of the nekton species considered in the analysis and the respective
129 categorisation in terms of EUFG, FMFG and fishery category is shown in Appendix (Table A.1).

130 **2.3. Environmental conditions**

131 In each study area a series of environmental variables was estimated using spatial data collected in previous
132 studies or measured *in situ* during this study. Variables not collected *in situ* were considered fixed in time (i.e.
133 temporal variability is assumed to be negligible over the study period), and are: % area covered by saltmarsh
134 (Mag. Acque, 2002; marsh); % area covered by seagrass bed habitat (*Cymodocea nodosa*, *Nanozostera noltii*
135 and *Zostera marina* monospecific and mixed assemblages; Rismondo et al., 2003; Curiel et al., 2014;
136 seagcover); bottom depth averaged by study area (in metres; Mag. Acque, 2002; bat); average distance of grid
137 cells composing each study area from the nearest sea inlet (metres; dist; evaluated on a regular grid with a cell
138 size of 100m); sand content in bottom sediments averaged by study area (as % sand; Mag. Acque - SELC,
139 2005; Mag. Acque - Thetis, 2005; sab); water residence time averaged by study area (days; Cucco et al., 2009;
140 restime); water speed averaged by study area (expressed as cm/s; Molinaroli et al., 2007; wsp). Variables
141 including temporal variability are: water salinity averaged by area in each sampling occasion (± 1 PSU –
142 Practical Salinity Units; this study; sal); water turbidity averaged by area and in each sampling occasion (± 1
143 FTU – Formazin Turbidity Units; this study; torb); water temperature recorded in each sampling occasion
144 ($\pm 0,1^\circ$ C; this study; temp); anomalies in water temperature ($^\circ$ C; Mag. Acque - SAMA, 2013; anoT).

145 **2.4. Anthropogenic pressures**

146 A set of indicators of anthropogenic pressure was developed, adapting the scheme proposed by Aubry and
147 Elliott (2006) according to the available knowledge for the Venice lagoon and to the relevance for nekton
148 fauna of each indicator (Table 1). Three main categories of pressures were identified, which were related to:
149 I) Morphology, II) Resource and Habitat Use and III) Environmental Quality.

150 Five indicators of Morphological pressure were selected (Table 1): intertidal area loss; seagrass habitat loss;
151 gross change in bathymetry; interference with hydraulic circulation due to the presence of human
152 infrastructures; relative sea level rise. The temporal variability of these indicators within the study period

153 (2001-2013) was assumed to be negligible and therefore only spatial variability (differences among areas) was
154 considered in the analysis. The Resource and Habitat Use category included five indicators (Table 1): artisanal
155 fishery; shellfish aquaculture; intensity of marina development; boat traffic; intensity of shipyards. The marina,
156 navigation and shipyard indicators were maintained fixed in time due to the lack of time series of data. Seven
157 indicators of Environmental Quality were selected (Table 1), representing: water chemical quality; sediment
158 chemical quality; sediment quality biological effects; ecological status of macrobenthos; chlorophyll-*a*
159 concentration; nutrients concentration; dissolved oxygen. Biological effects of sediment quality and benthic
160 state were kept fixed in time, due to lack of data available over time. When seasonal values were not available
161 for the other indicators, these were replaced with nearest-in-time values.
162 Values of each pressure indicator were standardised according to a five-level classification scoring (Table
163 A.2). Pressure levels were evaluated for each element of a 1km x 1km cell grid covering the lagoon extent and
164 the mean pressure level for each indicator in each study area was obtained by overlaying the grid maps

165 **2.5. Data Analysis**

166 **2.5.1. Model calibration**

167 Generalized Linear Models (GLMs; McCullagh and Nelder, 1989) were fitted to link response variables to
168 temporal, environmental and pressure predictors, as described in the next paragraphs. Models for number of
169 species (S) and biomass (B) of functional groups (univariate analyses) and for presence-absence and biomass
170 of species (multivariate analyses) were fitted considering different combinations of predictor variables and the
171 most suitable error term (depending on the type of response variable), in order to build the set of models
172 representing the different hypotheses considered in this study.

173 Response variables (univariate and multivariate analysis)

174 Nekton data were analysed both in a univariate and a multivariate fashion. Firstly, nekton data were
175 summarised in a set of variables -or metrics- quantifying different characteristics of the assemblage. These
176 metrics were computed as biomass and species richness of Estuarine Usage Functional Groups (16 metrics),
177 Feeding Mode Functional Groups (16 metrics) and fishery categories (6 metrics). All metrics were
178 independently used as response variables in GLMs fitted using Table 2 formulas (univariate analysis), and
179 choosing the most appropriate distribution family, after the visual inspection of the mean-variance relationship

180 (Warton, 2008; Warton et al., 2012). Secondly, following the approach of the `manyglm` software package
181 (Wang et al., 2012), binomial GLMs were fitted using presence/absence information of each species and
182 following the same formulation proposed for univariate data (Table 2), and inferences were carried out at the
183 assemblage level (Wang et al., 2012; Warton et al., 2012) (multivariate analysis). A similar approach was
184 replicated on biomass data, developing negative binomial GLMs for each species contributing to 95% of total
185 biomass and combining results in a global analysis.

186 Predictor variables

187 In order to represent the effects of the temporal variability on the response variables, the interaction between
188 the factors *Season* and *Year* was considered as a predictor in the model. Environmental conditions were
189 summarised by performing a principal component analysis, and the first three axes were considered as
190 predictors of environmental variability in fitted models. The predictors used as indices of pressures for the
191 three considered categories were the average values of the indicators within each category (Morphology
192 pressures, Resource and Habitat Use pressures - excluding the fishery -, Environmental Quality pressures). As
193 in this study a particular attention was paid to the role of the artisanal fishery, its indicator (see 2.4) was
194 considered as a separate predictor variable in model building.

195 Model structure

196 Models were fitted using different structures in order to hypothesise different contributions of the predictor
197 variables (Table 2). Eight model formulations, belonging to four model categories, were built addressing the
198 following hypotheses: none of the considered predictors affects the response variable (category *m0*); response
199 variable is influenced by temporal factor alone (category *m1*); response variable is affected by temporal and
200 environmental factors (category *m2*); response variable is affected by anthropogenic pressures (with or without
201 environmental conditions, and excluding fishery pressure) (category *m3.X*); and response variable is affected
202 by the fishery (with or without environmental conditions and other anthropogenic pressures) (category *m4.X*)
203 (Table 2). Not all models were nested (e.g. one included within another more complex one), but the category
204 of models were conceptually hierarchically designed (e.g. environmental variables or anthropogenic pressures
205 cannot be included in a model if temporal factor has not been considered yet).

206

207 **2.5.2. Comparison of models**

208 Fitted models were analysed to understand if nekton data showed a relationship with the anthropogenic
209 pressure categories, by:

210 - comparing models (both for univariate and multivariate approaches) by means of Likelihood Ratio tests
211 between chosen sets of nested models (Table 2). These comparisons were carried out to disentangle the
212 contribution of the different type of predictor variables considered (temporal factors, environmental conditions,
213 anthropogenic pressures and artisanal fishery pressure). For example, for a given metric, the comparison (Test
214 2.1, Table 2) of a model fitted using temporal factor and environmental conditions (m2.0, Table 2) with another
215 one including also anthropogenic pressures (excluding artisanal fishery) (m3.1, Table 2), suggests if the
216 inclusion of anthropogenic pressures significantly improves the model in the case that temporal factor and
217 environmental conditions were already included;

218 - computing the deviance explained by each model, in order to estimate the magnitude of the effect in addition
219 to its significance (see previous point);

220 - averaging all the candidate models (Table 1) for each univariate metric following an Information Theory
221 Criterion, to carry out global inference on the estimated averaged parameters for pressure variables (univariate
222 analysis). In particular, models were averaged using the AIC_c weights, and considering the 'top-models', i.e.
223 the models representing the 95% confidence model set (Grueber et al., 2011);

224 - averaging (unweighted mean values) the parameters of the models of all species for each category of pressure,
225 to summarise the effects of pressure indicators on nekton assemblage (multivariate analysis);

226 **3. Results**

227 **3.1. Nekton assemblage**

228 During the study period, 59 fish and two cephalopod species were recorded from fyke nets and included in the
229 analysis (Table A.1). Gobiidae, Sparidae, Syngnathidae and Mugilidae were the most numerous families in
230 terms of species numbers (with 8, 8, 7 and 5 species respectively), while *Sepia officinalis* and *Sepiola rondeletii*

231 were the only two species of cephalopod recorded. Two fish species, *Atherina boyeri* and *Zosterisessor*
232 *ophiocephalus*, accounted for 58% of the total biomass (39% and 19% respectively).

233 Nine species out of the 61 forming the whole nekton assemblage were targets of the fishery and accounted for
234 78% of the total biomass, with 11 incidental species accounting for an additional 16%. Thus, 94% of the total
235 biomass caught with fyke nets was composed by species with some commercial value (target + incidental
236 species), with the 41 discarded species representing only 6% of biomass of the analysed catches. Among
237 nekton species, eight EUFG were identified: anadromous (A), catadromous (C), estuarine (ES), solely
238 estuarine (ESs), freshwater stragglers (FS), marine estuarine-dependent (ME-D), marine estuarine-opportunist
239 (ME-O) and marine stragglers (MS). MS was the richest EUFG, with 26 species, followed by ME-O (12), ES
240 (11), ME-D (5) and ESs (4 species). A, C and FS EUFG were all represented by one species each. ES was the
241 dominant guild in terms of biomass (60% of the total), with *A. boyeri* and *Z. ophiocephalus* accounting for
242 95% of the guild biomass (64% and 31% respectively). ESs accounted only for 0.4% of the total biomass.

243 Eight FMFGs were identified: macrobenthivores (Bma), microbenthivores (Bmi), detritivores (DV),
244 hyperbenthivores/piscivores (HP), herbivores (HV), hyperbenthivores/zooplanktivores (HZ), omnivores (OV)
245 and planktivores (PL). Bmi and Bma were the FMFG with more species among nekton assemblage. *Z.*
246 *ophiocephalus*, *Solea solea* and *Platichthys flesus* accounted together for 73% of Bmi biomass, while *Z.*
247 *ophiocephalus*, *S. officinalis* and *P. flesus* accounted for 79% of Bma biomass. Twenty species were allocated
248 to the HP guild, with seven of them showing exclusively this feeding mode. *Z. ophiocephalus* and *S. officinalis*
249 were the dominant species within the HP guild, accounting for 75% in terms of biomass. Fourteen species were
250 HZ with only three of them being exclusive to this guild. *A. boyeri* alone accounted for 99% of the HZ guild
251 biomass. Six omnivorous species were found, all being allocated in multiple FMFG except for *Diplodus*
252 *puntazzo*. Mugilids accounted for the totality of the DV guild. Engraulids and clupeids entirely accounted for
253 the PL guild.

254 **3.2. Environmental conditions and anthropogenic pressures**

255 The five study areas were clearly ordered along the first principal component (Figure 2a), which can thus be
256 interpreted as a gradient from more dynamic and marine-like conditions, typically observed in proximity to
257 the sea inlets (areas CH and LD), towards more confined areas (LT and CZ). In addition, the observations
258 within each study area were scattered along the second principal component, due to the interannual and

259 seasonal variability (Figure 2a). The dispersion along the third component was similar to that of the second
260 axis with the points belonging to the different areas largely overlapping, but CH and LT - the least and the
261 most confined areas - indicated a lower turbidity level (Figure 2b). The first component of the PCA performed
262 on environmental variables identified a main spatial confinement gradient as ascribed to changes in water
263 residence time, seagrass cover, sediment grain size and water speed, while the second component was strongly
264 influenced by variables with a sharp seasonal variability (water temperature and dissolved oxygen) and the
265 third component was related to variables changing over time (water temperature anomaly) or in time and space
266 (turbidity) (Fig. 2b).

267 In the case of the PCA performed on anthropogenic pressures, it was not easy to recognise a geographical
268 pattern in the pressure gradient, as all three axes were influenced by variables representing pressures with a
269 strong spatial component (interference with hydrographic regime, relative sea level rise, gross change in
270 bathymetry) or variables changing over time both with a seasonal (e.g. average concentration of Dissolved
271 Inorganic Nitrogen) or inter-annual dynamic (area affected by aquaculture activities) (Fig. 2c and 2d).
272 However, the distribution of the areas along the first component partially corresponded to the one described
273 by the principal component analysis on environmental variables; with an ordination from the innermost area
274 (LT), where the loss of intertidal area, the sediment chemical quality and the macrobenthos state were the
275 greatest pressures, to the area closest to the sea inlet (CH), where pressures related to sea level rise, aquaculture,
276 interference with hydrographic regime pressures and the loss of seagrass were more important (Figure 2c).
277 However, on this ordination CZ differed in location compared to the previous PCA, as if pressures in CZ could
278 be associated with the ones of a less confined area (Figure 3a). The second axis combined LD and PL against
279 the others stations, due to a higher importance of pressures related to the intensity of navigation and density of
280 shipyards in these areas (Figure 2c), while the third component distinguished CZ and PL from the other areas
281 for being less affected by bathymetric change (Figure 2d).

282

283 **Figure 2 about here.**

284

3.3. Effects of temporal factors, environmental conditions, and anthropogenic pressures upon nekton assemblage

Univariate comparison

All the univariate GLMs for biomass-related metrics were fitted using a negative binomial distribution, while the models for the number of species metrics were fitted using a Poisson distribution. All the models explained a low to moderate proportion of deviance (Figure 3), with few exceptions related to metrics with very sparse (i.e. zero inflated) data matrix, probably due to overfitted models. All the metrics indicate a significant seasonal pattern, as the effect of temporal factor alone (test 0) was significant ($p < 0.001$) for all metrics, except for the biomass of catadromous EUFG, and for the species richness of anadromous and freshwater straggler EUFG and herbivore FMFG (Figure 3). It is worth noting that data for these groups of species were strongly zero inflated. In general, the metrics show a significant seasonal pattern.

The inclusion of environmental conditions (test 1) had a significant effect ($p < 0.01$) on all biomass metrics except for catadromous EUFG and omnivorous FMFG, and on the same species richness metrics for which temporal factor alone was significant. This means that for most of the metrics, the differences between observations can in part be explained by environmental conditions, in addition to the contribution of temporal factors.

When only temporal factor was already considered, the effect of including all anthropogenic pressures with the exception of the fishery (test 2.0) proved significant ($p < 0.05$) for most of the metrics (but see biomass of catadromous EUFG and species richness of anadromous EUFG and herbivores FMFG; Figure 3).

Including pressures with both temporal and environmental conditions already considered (test 2.1) significantly ($p < 0.05$) contributed to explain the variance of many metrics, with the exception of the biomass of incidental catches, catadromous, freshwater stragglers, marine estuary-dependent and detritivorous species and the number of species of target, total catches, anadromous EUFG and of herbivorous FMFG. Hence, in a large number of cases, adding the information on anthropogenic pressures to the temporal and environmental factors assists in explaining the variance in the nekton assemblage data.

In general, the inclusion of artisanal fishery pressure resulted in significance for a smaller number of cases (Figure 3). The inclusion of fishery pressures when temporal factors, environmental variables and the other

312 pressures were already included in models (test 3.2), resulted in significance of fewer metrics than in the case
313 of the inclusion of the fishery when temporal factor alone (test 3.1) or temporal factor and environmental
314 variables (test 3.0) were already included in the analysis. However, this is not true for all metrics: for those
315 metrics representing the biomass of the EUFG, significance was more likely for the fishery if all the other
316 factors were previously included in the model.

317

318 When considering the model averaged parameters associated to the different indicators of pressure (Figure 4),
319 the fishery showed a slightly negative effect only on biomass of discarded catches, anadromous and marine-
320 estuarine opportunist EUFGs and both detritivorous and planktivorous FMFGs. Similarly, the fishery
321 negatively affected the species number of incidental catches, estuarine and marine estuarine-opportunist
322 EUFGs and macrobenthivorous, detritivorous and hyperbenthivorous/piscivorous FMFGs. Overall though, the
323 fisheries effect on the considered metrics seemed to be negligible, since the coefficients associated with this
324 pressure were often very close to zero (Figure 4). Moreover, in many cases, these effects were characterised
325 by a small influence (low Akaike weights) on the averaged models. In addition, a positive effect of the fishery
326 was observed for biomass of target and total catches, of estuarine, solely estuarine and marine straggler EUFGs
327 and of benthivorous FMFGs (Figure 4).

328 Compared with the fishery, a stronger effect (larger β coefficients) of the other pressure categories was evident
329 (Figure 4). No common pattern could be identified among metrics, except for the biomass of the fisheries-
330 related metrics (discarded, incidental, target and total catches), which seemed to be negatively affected by both
331 morphology-, resource use- and environmental quality-related pressures. Overall, pressures related to
332 morphological alterations had marked negative effects on biomass of discarded, incidental, target and total
333 catches, as well as biomass of estuarine EUFG and of both microbenthivorous and omnivorous FMFGs.
334 Resource use pressures showed the most negative effects on biomass of discarded catches and
335 hyperbenthivorous/zooplanktivorous FMFG. Finally, environmental quality degradation had a particularly
336 negative effect on biomass of target and total catches, marine estuarine-opportunist EUFG and four FMFGs
337 (both macro- and microbenthivorous, hyperbenthivorous/psicivorous and planktivorous species).

338

339 **Figure 3 about here.**

340 **Figure 4 about here.**

341

342 ***Multivariate comparison***

343 Only the effect of temporal factor (test 0) was significant ($p < 0.001$) when analysing GLMs performed on
344 species presence/absence in the whole nekton assemblage (Table 3). None of the pressure categories proved
345 significant in determining species composition of the assemblage as a whole.

346 In contrast, a higher number of significant effects ($p < 0.01$) were detected when comparing GLMs performed
347 on the biomass distribution across the species in the assemblage. The effect of temporal factors (test 0) was
348 significant, as well as the effect of adding environmental conditions when temporal factors were already
349 considered (test 1), adding anthropogenic pressures excluding the fishery when temporal factors were already
350 considered (test 2.0), adding anthropogenic pressures excluding the fishery when both temporal factor and
351 environmental conditions were already considered (test 2.1) and adding fishery pressure when both temporal
352 factor and environmental conditions were already considered (test 3.0) (Table 3).

353 Five target species commonly found in fyke nets showed significant effects of anthropogenic pressures most
354 frequently across tests (see supplementary materials, Table A.3). *S. officinalis* showed a significant effect in
355 six tests (five of which performed on species presence/absence and one on biomass), while *A. boyeri* and *Liza*
356 *aurata* showed a significant effect in five tests (three of which performed on species presence/absence and two
357 on biomass). In addition, the effect of pressures also proved significant for the presence of *Pomatoschistus*
358 *minutus* and *Z. ophiocephalus* (three and two tests respectively). With both temporal and environmental factors
359 already considered (Test 2.1), pressures related to morphological alterations and resource use negatively
360 affected the probability of target species presence as *Anguilla anguilla*, *L. aurata*, *Sparus aurata* and *Z.*
361 *ophiocephalus*, with *A. boyeri* showing a negative effect only for resource use pressures. In contrast, the
362 presence of *P. flesus* and *P. minutus* (target species) was affected positively or not affected by both these
363 pressure categories. Similarly, environmental degradation had a negative effect on the presence of *P. flesus*,
364 *P. minutus*, *S. aurata* and *S. officinalis* with both temporal and environmental factors already considered, while
365 it showed a positive effect on *A. anguilla*, *A. boyeri*, *L. aurata* and *Z. ophiocephalus*. The fishery affected the
366 likelihood of four target species presence when it was considered as the only pressure (Test 3), having a
367 positive effect on *A. boyeri*, *L. aurata* and *Mugil cephalus* and a negative impact on *S. officinalis*.

368 When considered together with other pressure category (Test 3.2), in contrast, the fishery negatively affected
369 the presence of *A. anguilla* and *S. Officinalis* only, while being negligible for other target species. Additionally,
370 test 2.1 indicated that morphological pressures had a negative effect upon biomass of *A. boyeri* and *L. aurata*,
371 while both resource use and environmental quality pressures were positively related to biomass of these
372 species.

373 On average, pressures related to morphological changes, resource and habitat use, as well as pressures on
374 environmental quality showed negative effects on the biomass of the assemblage, while only pressures on
375 environmental quality negatively affected the average probability of presence of the single species within the
376 assemblage (Figure 5). Both for presence/absence and for biomass, the effect of fishery pressure seemed to be
377 negligible (Figure 5).

378

379 **Figure 5 about here.**

380

381 **4. Discussion**

382 We investigated the potential use of data gathered during the monitoring of the artisanal fishery in the Venice
383 lagoon to describe the relationship between nekton assemblage (including fish and cephalopods) and
384 anthropogenic pressures in transitional waters.

385 The Venice lagoon represents a good case study, since a traditional form of fishing is carried out within its
386 boundaries by local fishermen throughout the year in most part of the basin (Provincia di Venezia, 2009;
387 Pranovi et al., 2013). Moreover, this lagoon, given its wide area and complex hydro-morphology, is
388 characterised by a mosaic of habitats and multiple environmental gradients that lead to a high level of
389 environmental variability (Solidoro et al., 2010). The Venice lagoon is also subjected to a variety of pressures,
390 and their variability in space and time is rather well documented (Franco et al., 2009a; Solidoro et al., 2009).
391 The combination of natural heterogeneity in time and space with the unevenness of anthropogenic pressures
392 and impacts can change ecosystem organisation and functioning (Brigolin et al., 2014). Therefore it was crucial
393 to take into account these sources of heterogeneity, evaluating anthropogenic pressures, environmental
394 conditions and nekton assemblage in different areas, different times of the year, and between years. For the

395 areas considered in this study, spatial variability in environmental conditions and anthropogenic pressures
396 seems to be stronger than the temporal, as highlighted by the Principal Component Analysis (Figure 2). This
397 is due to the marked spatial variability, but also due to the assumptions at the basis of this work (pressures
398 definition), and the data availability constrains.

399 The main anthropogenic pressure gradients within the lagoon are associated with morphological changes such
400 as habitat loss and alteration of the hydrodynamic conditions, as previously noted (Franco et al., 2009a;
401 Molinaroli et al., 2009; Sarretta et al., 2010), with also alterations of environmental quality being an important
402 pressure category. Indeed, despite the stricter regulations and the decline of industrial activities in the last
403 decades, persistent contaminants such as heavy metals and organic compounds are still stored in lagoon
404 sediments, directly affecting the associated benthic compartment (Secco et al., 2005; Bernardello et al., 2006).

405 The whole nekton assemblage as assessed by using fyke net catches did not show a significant relationship
406 with the environmental variability of the lagoon when considering its species composition, whereas the
407 biomass structure of assemblages indicated the effect of temporal, environmental and pressure factors. Indeed,
408 species occurrence in the catches changed significantly according only to season and year (test 0), as a result
409 of the temporal dynamics in nekton populations such as recruitment and migrations (Elliott and Hemingway,
410 2002). Environmental variability and anthropogenic pressure did not have significant effects on catch
411 composition, in terms of species richness and presence-absence of species.

412 On the contrary, anthropogenic pressures significantly explained part of the variability on the biomass of
413 catches, in addition to both temporal variability alone (test 2.0) and with environmental conditions (test 2.1).
414 Species biomass in the assemblage proved to be sensitive not only to pressures related with morphological
415 change, resource use (excluding artisanal fishery) and environmental quality, but also in a smaller degree to
416 the effects of the artisanal fishery. The latter may be due to changes in population structure related to the
417 removal of individuals of target species, hence leading to the observed alteration in the biomass of the
418 assemblage. Regarding the effects of morphological degradation, it is not easy to identify the effects of physical
419 disturbance of the environment for mobile or migratory species, as they may change their distribution and
420 behaviour, but this could be simpler for resident species (Marchand et al., 2002). Some authors have suggested
421 that the degradation or loss of habitats affects fish species richness in estuaries (Harrison and Whitfield, 2004;
422 Cabral et al., 2012; Harrison and Kelly, 2013). Our results confirm that the effects of pressures acting on the

423 lagoon morphology are stronger for resident species (e.g. ES biomass and biomass of *Z. ophiocephalus* and *S.*
424 *officinalis* see fig. 4 and Tab A.3), but they can be significant also for migrant species (e.g. ME-D biomass and
425 biomass of *A. anguilla* and *L. saliens*; see fig. 4 and Tab A.3). However, it is interesting to note that these
426 effects can be manifested in our study as impacts on biomasses (univariate and multivariate tests) rather than
427 on metrics accounting for species richness and assemblage composition (presence/absence tests). Many studies
428 (e.g. Brandão et al., 2013; Fonseca et al., 2014; Gonçalves et al., 2014) describe the effects of chemical
429 pollutants on fishes, but these effects usually cannot be easily observed and quantified at the population level,
430 especially if pollution is present at sub-lethal levels (Hamilton et al., 2015). Marchand et al. (2002) suggest
431 that for some types of pollutants, the effects are unlikely to be detrimental, because fishes avoid highly polluted
432 areas. The significant effects of pressures on quality of matrices observed in this study are probably more
433 related to enrichment in nutrients and organic matter, leading to anoxic crises, than to chemical pollution (in
434 agreement with the patterns observed, for example, by Uriarte and Borja, 2009).

435 Several of the metrics responded in a coherent way to the impacts of human pressures. As an example, biomass
436 of estuarine resident species, benthivores, hyperbenthivores/piscivores, target species and of total catches
437 indicated a clearly negative response to morphological alterations. This result could be related to the role within
438 these categories of some species, for example the grass goby *Z. ophiocephalus*, for which the impacts on the
439 essential habitat for its reproduction (i.e. seagrass meadows; Malavasi et al., 2004b) might determine a decrease
440 on the population biomass. However, biomass of both target and total catches demonstrated a strong negative
441 response to all three main pressure categories.

442 In this study, the biomass of marine estuarine-opportunists was negatively influenced by pressures acting on
443 quality of matrices. Similar results were found by Amara et al., (2009), who recorded lower growth and body
444 condition in juveniles of *P. flesus* (marine migrant) exposed to higher levels of chemical pollution in French
445 estuaries. On the other hand, the biomass of solely estuarine species revealed a positive relationship with
446 pressures related to environmental quality in the Venice lagoon. The species of this guild (e.g. the lagoon goby
447 *Knipowitshia panizzeae*) are more likely to be adapted to high levels of natural variability typical of transitional
448 waters (Franco et al., 2008), and therefore might be also able to cope with poor environmental conditions.
449 Indeed, some studies suggest that estuarine resident species may be able to develop resistance to chemical
450 pollution as an adaptation to long-term exposure (Nacci et al., 1999; Matthiessen and Law, 2002; Fonseca et

451 al., 2013) and this might give them an additional competitive advantage in colonising and benefiting from
452 lower quality areas, where less tolerant species are excluded or of reduced abundance.

453 The partial differences in the response of different guild categories suggest a certain degree of complementarity
454 between them. Hence, this supports the value of the followed approach in interpreting nekton assemblage
455 structure, due to the use of functional rather than taxonomical categories (Elliott et al., 2007; Franco et al.,
456 2008; Mouillot et al., 2013; Henriques et al., 2014), suggesting the implementation of an indicator set (i.e.
457 metrics) of different aspects of the community potentially influenced by anthropogenic impacts.

458 **4.1. Artisanal fishery**

459 The role of the artisanal fishery in affecting the nekton assemblage was considered in conjunction with other
460 anthropogenic pressures, as suggested by Blaber et al. (2000). Overall, the fyke net fishery was less important
461 compared with other pressure categories. For biomass and species composition of most EUFG, FMFG and of
462 the whole assemblage, the effect of the artisanal fishery was negligible (univariate analysis, Figures 3 and 4).
463 A marginal positive response to the pressure was observed on biomass of estuarine and solely estuarine
464 residents, as well as on biomass of microbenthivores (a FMFG including a large variety of species belonging
465 to different EUFG and fishery categories). Remarkably, fishery pressure indicated a similar positive effect
466 upon biomass of both target and total catches, although with a high associated variability.

467 Fishing in the Venice lagoon is performed using a variety of traditional gears, with fyke nets being the most
468 important (Granzotto et al., 2001; Provincia di Venezia, 2009). The extraction of data from this activity is
469 similar in many ways to a scientific survey: it is carried out throughout the year and the basin using a single
470 type of gear, which has similar length, height and mesh size and is normally deployed for a comparable amount
471 of time (Provincia di Venezia, 2009; Pranovi et al., 2013). All these characteristics allow the spatial and
472 temporal comparison of collected samples by means of a standardised CPUE. Nevertheless, selection of fishing
473 sites is mainly made in order to maximise yields. This may explain both the positive effect on target species
474 and the negligible effect of the fishery in other cases, since a higher fishing effort is often justified in areas
475 where higher abundance and biomass of targeted species are known to be supported. However, this positive
476 link can persist in time only if the exploitation is carried out at biological sustainable levels. Indeed, the overall
477 effort of the artisanal fishery (average of the whole basin) was relatively stable over the study period, even if
478 it demonstrates a strong seasonality and high spatial variability (Provincia di Venezia, 2015). The present

479 levels of fishing activities are lower than the ones recorded in the past, and indications suggest that changes in
480 yields recorded in the last decades are not caused by overfishing, but rather by other human-induced changes
481 in environment quality (Libralato et al., 2004).

482 The results highlighted above suggest that the fyke net-based artisanal fishery in the Venice lagoon is possibly
483 undertaken at a sustainable level for the nekton fauna, considering that no significant effect was observed on
484 the species (or group of species) composition or biomass of the lagoon nekton assemblages. Furthermore
485 discarded levels are low, with up to 6% of the total biomass being discarded (Pranovi et al., 2013). However
486 it should be noted that benthic invertebrate species, not considered in this study, can also be caught by fyke
487 nets, and can even represent a significant share of the total biomass in the catch, both in terms of target species
488 (i.e. crabs; Pranovi et al., 2013) and discarded species (e.g. gastropods). Only *ad hoc* fishery-independent
489 surveys could fully investigate the overall sustainability of this type of artisanal fishery further. Such surveys
490 would need to take into account the relationship between fishing effort and the response of the ecosystem,
491 which is not necessarily linear. The effects of fishing effort could therefore be only evident after a certain
492 critical threshold has been reached (Henriques et al., 2014). It was important however, for the goals of the
493 present work, to understand if the activity used to gather information on nekton assemblage negatively affected
494 the assemblage itself. Fisheries in transitional waters are poorly studied, despite the fact that they represent a
495 relevant economic activity in these ecosystems (Pérez-Ruzafa and Marcos, 2012). In particular, while many of
496 the existing studies focus on the assessment of fisheries within transitional waters in terms of its impact on fish
497 communities and overall ecological quality (Blaber et al., 2000; Rodríguez-Climent et al., 2012; Guillemot et
498 al., 2014), there has not been any research regarding the potential use of fishery data to assess ecological status,
499 even if it is clear that yields are strongly influenced by environmental conditions and anthropogenic pressures
500 (Pérez-Ruzafa and Marcos, 2012).

501 In Europe, appropriately modified fishing gears are widely employed in scientific monitoring of fish fauna in
502 transitional waters. Fine mesh, small beach seine nets are used in shallow water habitats within Italian coastal
503 lagoons (Franco et al., 2006, 2009b), with modified beach seine nets being used also for the WFD assessment
504 of transitional waters in the UK, where a multi-gear approach (i.e. including also the use of fyke nets, beam
505 trawls and otter trawls) is applied (WFD-UKTAG, 2009). In French Mediterranean lagoons, fyke nets similar
506 to those used by local fishermen are preferred (Mouillot et al., 2005; Brehmer et al., 2013). Beam and otter

507 trawls are widely used in deeper habitats across estuaries in northern Europe, such as in UK and Germany
508 (Elliott and Hemingway, 2002), and are also employed in French and Portuguese Atlantic estuaries (Pasquaud
509 et al., 2010; Fonseca et al., 2013).

510 All sampling methods are affected by a degree of selectivity and their catch efficiency depends on both target
511 species and habitat characteristics. Hence, due to the spatial diversity of transitional water ecosystems and the
512 different fish assemblages they support, the choice of sampling gear is critical (Rozas and Minello, 1997;
513 Elliott and Hemingway, 2002). Franco et al. (2012) showed that active gears (i.e. seines) gather punctual
514 information at small spatial and temporal scales, while passive gears (i.e. fyke nets) tend to integrate on larger
515 time (e.g. day-night cycle) and space scales, without detecting eventual different roles of habitats. In this light
516 fyke nets could be more suitable for the comparison of relatively large areas, such as the ones considered here
517 (14 - 39 km²), or for the assessment of whole water bodies, where multi-gear approach is unfeasible.

518 **4.2. Implications for the evaluation of ecological status**

519 Our study supports the hypothesis that artisanal fishery data could represent an effective source for ecological
520 status assessment of the Venice lagoon, since it provides a non-invasive sampling method and allows the use
521 of pressure-sensitive metrics. Several fish-based multi-metric indices have been developed in recent years to
522 assess the environmental status of European transitional waters under the Water Framework Directive (e.g.
523 Coates et al., 2007; Franco et al., 2009b; Breine et al., 2010; Delpech et al., 2010; Cabral et al., 2011). As
524 shown by Pérez-Domínguez et al. (2012), most of the metrics included in such indices are measures of
525 community composition and structure, such as species richness and diversity. In addition, many of these
526 indices follow the guild approach (Elliott and Dewailly, 1995; Elliott et al., 2007; Pérez-Domínguez et al.,
527 2012). In contrast, biomass does not seem to be incorporated into published fish-based indices complying with
528 WFD (Cabral et al., 2012; Pérez-Domínguez et al., 2012), even if a few proposed methods include fyke net
529 sampling to obtain fish data (Coates et al., 2007; Breine et al., 2010). In fact, biomass could serve as a proxy
530 indicator for the secondary production of a system, which can be affected by loss or degradation of transitional
531 habitats (Deegan et al., 1997; Hughes et al., 2002).

532 Our study indicates that metrics calculated on biomass are more effective in detecting anthropogenic stressors
533 compared with measures of species composition, when using fyke nets as sampling gear. In the absence of size
534 related metrics, biomass also proved to be a good indicator of the effects of fishing activities, as recorded in

535 other aquatic environments (Vallès and Oxenford, 2014). Compared to biomass, the use of species
536 presence/absence led to the coexistence of a variety of different responses (highly positive, negative and
537 negligible) in our study, ultimately resulting in an ambiguous representation. The importance of the
538 interpretability of response variables, their sensitivity to anthropogenic pressures and robustness against
539 natural ‘noise’ is crucial when evaluating ecological status (Rice, 2003; Noges et al., 2009).

540 The use of biomass-based metrics to detect changes in nekton communities was previously proposed by other
541 authors (Guillemot et al., 2014; Henriques et al., 2014; Vallès and Oxenford, 2014). Here, we suggest that an
542 evaluation of ecological status based on fishery monitoring data should focus on biomass data grouped
543 considering a functional guild approach. This work shows some metrics responding to anthropogenic
544 pressures, but further considerations are needed to choose which metrics, or combination of metrics, would
545 result in the most effective index. Here a wide range of characteristics of nekton assemblage were analysed in
546 relation to anthropogenic pressures and it was not possible to analyse in detail the interpretation of each
547 comparison, nor to examine the causal link with human activities or to search for a combination of metrics
548 maximising the correlation with anthropogenic pressures. In fact, even if the objective is to develop an index
549 with a strong responsiveness to human disturbance, if only information on bivariate correlations with pressures
550 are available, it is not possible to say which combination will result in a sensitive multi-metric index
551 (Schoolmaster et al., 2012). The selected method for data analysis was based on the building of a series of
552 nested models for the observed data, and the subsequent comparison of chosen sets of such models. This
553 allowed us to explicitly answer ecological questions (Warton et al., 2014) based on alternative *a priori*
554 hypotheses about the different contribution of seasonal and annual variability, environmental heterogeneity
555 and anthropogenic pressures in affecting the nekton assemblage. It was possible to test the effect of pressures
556 on the nekton assemblage as a whole and on a series of macro-descriptors summarising important
557 characteristics of the assemblage. This model-based approach is acknowledged to be a more interpretable,
558 flexible and efficient way to handle ecological data, compared to other classical multivariate analysis
559 techniques (Warton et al., 2014). This approach allowed us to disentangle the contribution the different types
560 of predictor variables, ultimately enabling the discrimination between the effects of anthropogenic pressures
561 and environmental variability on nekton assemblage. We argue that such an approach represents an effective
562 way to cope with the “estuarine quality paradox”, i.e. the difficulty to detect anthropogenically-induced stress

563 in estuaries due to the adaptation of the biota to the naturally high variability within such ecosystems (Elliott
564 and Quintino, 2007; Elliott and Whitfield, 2011). Since biological indicators must be able to measure the
565 ‘signal’ of anthropogenic effects over the ‘noise’ of this natural variability (Whitfield and Elliott, 2002), the
566 estuarine quality paradox is regarded as a strong handicap to assess the ecological status of transitional waters
567 (Dauvin and Ruellet, 2009).

568 **5. Conclusions**

569 The results of this study have major implications on several aspects of research and management of transitional
570 water ecosystems: (i) indicating the critical importance of evaluating the relationship between nekton
571 assemblage and anthropogenic pressures, and suggesting the most suitable metrics for this purpose considering
572 the estuarine quality paradox (Elliott and Quintino, 2007); (ii) confirming the effectiveness of the guild
573 approach in such a context, particularly by considering biomass-related metrics; (iii) exemplifying the
574 effectiveness of model based community analysis, which allows to explicit answers to ecological questions
575 and to test *a priori* formulated hypotheses; (iv) investigating a poorly researched subject in estuarine ecology
576 such as the role of the artisanal fishery in affecting nekton communities, and disentangling the contribution of
577 this activity from other anthropogenic pressures; (v) suggesting that an artisanal fishery performed with fyke
578 nets can represent a viable source of ecological data in coastal lagoons; (vi) suggesting that fyke net-based
579 artisanal fishery in the Venice lagoon, carried out at the present level of effort, is possibly sustainable with
580 regard to the nekton assemblage, since its role as a pressure is negligible compared with the other
581 anthropogenic stressors.

582

583

6. Acknowledgements

584 This study has been partially funded by the Italian Ministry of Universities and Research (PRIN grants
585 2009W2395), by Corila (Consorzio Ricerche Lagunari), and by the local water authority "Provveditorato
586 Interregionale alle OO. PP. del Veneto- Trentino Alto Adige – Friuli Venezia Giulia". We thank the
587 Provveditorato Interregionale alle OO. PP. del Veneto- Trentino Alto Adige – Friuli Venezia Giulia for
588 providing data for the evaluation of anthropogenic pressures, Riccardo Fiorin and Federico Riccato (Laguna
589 Project s.n.c.) for carrying out the field work. We are grateful also to the fishermen Ivan Bognolo, Umberto
590 Costantini and Mario Salvagno for their collaboration in fieldwork activities. We also thank two anonymous
591 reviewers for their helpful comments and Luise Roberts for the English revision of the text.

592

7. References

- 594 Alves, D.M., Cristo, M., Sendão, J., Borges, T.C., 2006. Diet of the cuttlefish *Sepia officinalis* (Cephalopoda:
595 Sepiidae) off the south coast of Portugal (eastern Algarve). *J. Mar. Biol. Assoc. UK* 86, 429.
596 doi:10.1017/S0025315406013312
- 597 Amara, R., Selleslagh, J., Billon, G., Minier, C., 2009. Growth and condition of 0-group European flounder,
598 *Platichthys flesus* as indicator of estuarine habitat quality. *Hydrobiologia* 627, 87–98.
599 doi:10.1007/s10750-009-9717-9
- 600 Apitz, S., Barbanti, A., Bernstein, A.G., Bocci, M., Delaney, E., Montobbio, L., 2007. The assessment of
601 sediment screening risk in Venice Lagoon and other coastal areas using international sediment quality
602 guidelines. *J. Soils Sediments* 7, 326–341. doi:http://dx.doi.org/10.1065/jss2007.08.246
- 603 Ardizzone, G.D., Cataudella, S., Rossi, R., 1988. Management of coastal lagoon fisheries and aquaculture in
604 Italy. *AO Fish. Tech. Pap.* 293, 1–103.
- 605 ARPAV-ISPRA, 2013. Monitoraggio della laguna di Venezia ai sensi della Direttiva 2000/60/CE. Valutazione
606 dei dati acquisiti nel monitoraggio ecologico 2011-2012 ai fini della classificazione ecologica dei corpi
607 idrici lagunari (elementi di qualità fisico-chimica e chimici, .
- 608 Aubry, A., Elliott, M., 2006. The use of environmental integrative indicators to assess seabed disturbance in
609 estuaries and coasts: Application to the Humber Estuary, UK. *Mar. Pollut. Bull.* 53, 175–185.
- 610 Bernardello, M., Secco, T., Pellizzato, F., Chinellato, M., Sfriso, A., Pavoni, B., 2006. The changing state of
611 contamination in the Lagoon of Venice. Part 2: heavy metals. *Chemosphere* 64, 1334–45.
612 doi:10.1016/j.chemosphere.2005.12.033
- 613 Blaber, S., Cyrus, D.P., Albaret, J.-J., Chong Ving Ching, Day, J.W., Elliott, M., Fonseca, M.S., Hoss, D.E.,
614 Orensanz, J., Potter, I.C., Silvert, W., 2000. Effects of fishing on the structure and functioning of estuarine
615 and nearshore ecosystems. *ICES J. Mar. Sci.* 57, 590–602. doi:10.1006/jmsc.2000.0723
- 616 Blanc, A., Daguzan, J., 1998. Artificial surfaces for cuttlefish eggs (*Sepia officinalis* L.) in Morbihan Bay,
617 France. *Fish.Res.* 38, 225–231.
- 618 Brandão, F., Correia, A.T., Gonçalves, F., Nunes, B., 2013. Effects of anthropogenic metallic contamination
619 on cholinesterases of *Gambusia holbrooki*. *Mar. Pollut. Bull.* 76, 72–76.
620 doi:10.1016/j.marpolbul.2013.09.029
- 621 Brehmer, P., Laugier, T., Kantoussan, J., Galgani, F., Mouillot, D., 2013. Does coastal lagoon habitat quality
622 affect fish growth rate and their recruitment? Insights from fishing and acoustic surveys. *Estuar. Coast.
623 Shelf Sci.* 126, 1–6. doi:10.1016/j.ecss.2013.03.011
- 624 Breine, J., Quataert, P., Stevens, M., Ollevier, F., Volckaert, F.A.M., 2010. A zone-specific fish-based biotic
625 index as a management tool for the Zeeschelde estuary (Belgium). *Mar. Pollut. Bull.* 60, 1099–1112.
- 626 Brigolin, D., Facca, C., Franco, A., Franzoi, P., Pastres, R., Sfriso, A., Sigovini, M., Soldatini, C., Tagliapietra,
627 D., Torricelli, P., Zucchetta, M., Pranovi, F., 2014. Linking food web functioning and habitat diversity
628 for an ecosystem based management: A Mediterranean lagoon case-study. *Mar. Environ. Res.* 97, 58–
629 66. doi:10.1016/j.marenvres.2014.02.006
- 630 Cabral, H.N., Fonseca, V.F., Gamito, R., Goncalves, C.I., Costa, J.L., Erzini, K., Goncalves, J., Martins, J., Leite,
631 L., Andrade, J.P., Ramos, S., Bordalo, A., Amorim, E., Neto, J.M., Marques, J.C., Rebelo, J.E., Silva, C.,

- 632 Castro, N., Almeida, P.R., Domingos, I., Gordo, L.S., Costa, M.J., 2012. Ecological quality assessment
633 of transitional waters based on fish assemblages in Portuguese estuaries: The Estuarine Fish Assessment
634 Index (EFAI). *Ecol. Indic.* 19, 144–153. doi:10.1016/j.ecolind.2011.08.005
- 635 Caniglia, G., Borella, S., Curiel, D., Nascimbeni, P., Paloschi, A.F., Rismondo, A., Scarton, F., Tagliapietra,
636 D., Zanella, L., 1992. Distribuzione delle fanerogame marine [*Zostera marina* L., *Zostera noltii* Hornem,
637 *Cymodocea nodosa* (Ucria) Asch.] in laguna di Venezia. *Lav. - Soc.Ven.Sc.Nat.* 17, 137–150.
- 638 Carbognin, L., Teatini, P., Tomasin, A., Tosi, L., 2010. Global change and relative sea level rise at Venice:
639 What impact in term of flooding. *Clim. Dyn.* 35, 1055–1063. doi:10.1007/s00382-009-0617-5
- 640 Cataudella, S., Ferlin, P., 1984. Cataudella S., Ferlin P. (1984). Aspects de basse technologie dans
641 l'aménagement des ressources piscicoles et le développement de l'aquaculture dans les lagunes. *Stud.*
642 *Rev. GFCM/Etud. Rev. CGPM* 2, 567–591.
- 643 Chauvet, C., 1988. Manuel sur l'aménagement des pêches dans lagunes côtières: la bordigue méditerranéenne.
644 FAO Doc. Tech. Pêches 290, 77 p.
- 645 Coates, S., Waugh, A., Anwar, A., Robson, M., 2007. Efficacy of a multi-metric fish index as an analysis tool
646 for the transitional fish component of the Water Framework Directive. *Mar. Pollut. Bull.* 55, 225–240.
647 doi:10.1016/j.marpolbul.2006.08.029
- 648 Costanza, R., Arge, R., de Groot, R., Farberk, S., Grasso, M., Hannon, B., Limburg, K., Naeem, S., O'Neill,
649 R.V., Paruelo, J., Raskin, R.G., Suttonkk, P., 1997. The value of the world's ecosystem services and
650 natural capital. *Nature* 387, 253–260.
- 651 Cucco, A., Umgiesser, G., Ferrarin, C., Perilli, A., Canu, D.M., Solidoro, C., 2009. Eulerian and lagrangian
652 transport time scales of a tidal active coastal basin. *Ecol. Modell.* In Press, , 913–922.
653 doi:10.1016/j.ecolmodel.2009.01.008
- 654 Curiel, D., Checchin, E., Miotti, C., Pierini, A., Rismondo, A., 2014. Praterie a fanerogame marine della laguna
655 di Venezia - aggiornamento cartografico al 2010 e confronto storico. *Lav. Soc. Ven. Sc. Nat.* 39, 55–66.
- 656 D'Alpaos, L., 2010. Fatti e misfatti di idraulica lagunare. La laguna di Venezia dalla diversione dei fiumi alle
657 nuove opere delle bocche di porto. Istituto Veneto di Scienze Lettere ed Arti., Venice.
- 658 Dauvin, J.-C., Ruellet, T., 2009. The estuarine quality paradox: Is it possible to define an ecological quality
659 status for specific modified and naturally stressed estuarine ecosystems? *Mar. Pollut. Bull.* 59, 38–47.
660 doi:10.1016/j.marpolbul.2008.11.008
- 661 Deegan, L.A., Finn, J.T., Ayvazian, S.G., Ryder-Kieffer, C.A., Buonaccorsi, J., 1997. Development and
662 validation of an Estuarine Biotic Integrity Index. *Estuaries* 20, 601–617.
- 663 Delpech, C., Courrat, A., Pasquaud, S., Lobry, J., Le Pape, O., Nicolas, D., Boët, P., Girardin, M., Lepage, M.,
664 2010. Development of a fish-based index to assess the ecological quality of transitional waters: the case
665 of French estuaries. *Mar. Pollut. Bull.* 60, 908–18. doi:10.1016/j.marpolbul.2010.01.001
- 666 Elliott, M., 2002. The role of the DPSIR approach and conceptual models in marine environmental
667 management: an example for offshore wind power. *Mar.Poll.Bull.* 44, iii–vii.
- 668 Elliott, M., Dewailly, F., 1995. The structure and components of European estuarine fish assemblages.
669 *Neth.J.Aquat.Ecol.* 29, 397–417.
- 670 Elliott, M., Hemingway, K.L., 2002. *Fishes in Estuaries*. Blackwell Science, Oxford.

- 671 Elliott, M., Quintino, V., 2007. The Estuarine Quality Paradox, Environmental Homeostasis and the difficulty
672 of detecting anthropogenic stress in naturally stressed areas. *Mar. Pollut. Bull.* 54, 640–645.
673 doi:10.1016/j.marpolbul.2007.02.003
- 674 Elliott, M., Whitfield, A.K., Potter, I.C., Blaber, S.J.M., Cyrus, D.P., Nordlie, F.G., Harrison, T.D., 2007. The
675 guild approach to categorizing estuarine fish assemblages: A global review. *Fish Fish.* 8, 241–268.
676 doi:10.1111/j.1467-2679.2007.00253.x
- 677 Fonseca, V.F., Vasconcelos, R.P., França, S., Serafim, a., Lopes, B., Company, R., Bebianno, M.J., Costa,
678 M.J., Cabral, H.N., 2014. Modeling fish biological responses to contaminants and natural variability in
679 estuaries. *Mar. Environ. Res.* 96, 45–55. doi:10.1016/j.marenvres.2013.10.011
- 680 Fonseca, V.F., Vasconcelos, R.P., Gamito, R., Pasquaud, S., Gonçalves, C.I., Costa, J.L., Costa, M.J., Cabral,
681 H.N., 2013. Fish community-based measures of estuarine ecological quality and pressure–impact
682 relationships. *Estuar. Coast. Shelf Sci.* 134, 128–137. doi:10.1016/j.ecss.2013.02.001
- 683 Franco, A., Elliott, M., Franzoi, P., Torricelli, P., 2008. Life strategies of fishes in European estuaries: the
684 functional guild approach. *Mar. Ecol. Prog. Ser.* 354, 219–228. doi:10.3354/meps07203
- 685 Franco, A., Franzoi, P., Malavasi, S., Riccato, F., Torricelli, P., 2006. Fish assemblages in different shallow
686 water habitats of the Venice Lagoon. *Hydrobiologia* 555, 159–174.
- 687 Franco, A., Pérez-Ruzafa, A., Drouineau, H., Franzoi, P., Koutrakis, E.T., Lepage, M., Verdiell-Cubedo, D.,
688 Bouchoucha, M., López-Capel, A., Riccato, F., Sapounidis, A., Marcos, C., Oliva-Paterna, F.J., Torralva-
689 Forero, M., Torricelli, P., 2012. Assessment of fish assemblages in coastal lagoon habitats: Effect of
690 sampling method. *Estuar. Coast. Shelf Sci.* 112, 115–125. doi:10.1016/j.ecss.2011.08.015
- 691 Franco, A., Riccato, F., Torricelli, P., Franzoi, P., 2009a. Fish assemblage response pressures in the Venice
692 lagoon to environmental. *Transitional Waters Bull.* 3, 29–44. doi:10.1285/i1825229Xv
- 693 Franco, A., Torricelli, P., Franzoi, P., 2009b. A habitat-specific fish-based approach to assess the ecological
694 status of Mediterranean coastal lagoons. *Mar. Pollut. Bull.* 58, 1704–1717.
695 doi:10.1016/j.marpolbul.2009.06.016
- 696 Franzoi, P., Franco, A., Torricelli, P., 2010. Fish assemblage diversity and dynamics in the Venice lagoon.
697 *Rend. Lincei* 21, 269–281. doi:10.1007/s12210-010-0079-z
- 698 Froese, R., Pauly, D., 2015. FishBase. World Wide Web electronic publication. from <http://www.fishbase.org>.
- 699 Gonçalves, C., Martins, M., Diniz, M.S., Costa, M.H., Caeiro, S., Costa, P.M., 2014. May sediment
700 contamination be xenoestrogenic to benthic fish? A case study with *Solea senegalensis*. *Mar. Environ.*
701 *Res.* 99, 170–178. doi:10.1016/j.marenvres.2014.04.012
- 702 Granzotto, A., Franzoi, P., Longo, A., Pranovi, F., Torricelli, P., 2001. La pesca nella laguna di Venezia : un
703 percorso di sostenibilità nel recupero delle tradizioni Lo stato dell’arte. *Rapp. Sullo Svilupp. Sostenibile.*
704 *Fond. Eni Enrico Mattei.*
- 705 Granzotto, A., Libralato, S., Pranovi, F., Raichevich, S., Giovanardi, O., 2004. Comparison between artisanal
706 and industrial fisheries using ecosystem indicators. *Chem. Ecol.* 20, S43–449.
- 707 Grueber, C.E., Nakagawa, S., Laws, R.J., Jamieson, I.G., 2011. Multimodel inference in ecology and
708 evolution: challenges and solutions. *J. Evol. Biol.* 24, 699–711. doi:10.1111/j.1420-9101.2010.02210.x
- 709 Guillemot, N., Chabanet, P., Kulbicki, M., Vigliola, L., Léopold, M., Jollit, I., Le Pape, O., 2014. Effects of
710 fishing on fish assemblages in a coral reef ecosystem: From functional response to potential indicators.

- 711 Ecol. Indic. 43, 227–235. doi:10.1016/j.ecolind.2014.02.015
- 712 Hamilton, P.B., Cowx, I.G., Oleksiak, M.F., Griffiths, A.M., Grahn, M., Stevens, J.R., Carvalho, G.R., Nicol,
713 E., Tyler, C.R., 2015. Population-level consequences for wild fish exposed to sublethal concentrations of
714 chemicals - a critical review. *Fish Fish.* n/a–n/a. doi:10.1111/faf.12125
- 715 Harrison, T.D., Kelly, F.L., 2013. Development of an estuarine multi-metric fish index and its application to
716 Irish transitional waters. *Ecol. Indic.* 34, 494–506. doi:10.1016/j.ecolind.2013.06.018
- 717 Harrison, T.D., Whitfield, A.K., 2004. A multi-metric fish index to assess the environmental condition of
718 estuaries. *J.Fish Biol.* 65, 683–710.
- 719 Henriques, S., Pais, M.P., Vasconcelos, R.P., Murta, A., Azevedo, M., Costa, M.J., Cabral, H.N., 2014.
720 Structural and functional trends indicate fishing pressure on marine fish assemblages. *J. Appl. Ecol.* 51,
721 623–631. doi:10.1111/1365-2664.12235
- 722 Hering, D., Borja, A., Carstensen, J., Carvalho, L., Elliott, M., Feld, C.K., Heiskanen, A.-S., Johnson, R.K.,
723 Moe, J., Pont, D., Solheim, A.L., de Bund, W. Van, 2010. The European Water Framework Directive at
724 the age of 10: a critical review of the achievements with recommendations for the future. *Sci. Total*
725 *Environ.* 408, 4007–19. doi:10.1016/j.scitotenv.2010.05.031
- 726 Hughes, J., Deegan, L., Weaver, M., Costa, J., 2002. Regional application of an index of estuarine biotic
727 integrity based on fish communities. *Estuaries and Coasts* 25, 250–263.
- 728 Kapetsky, J.M., Lasserre, G. (Eds.), 1984. Management of coastal lagoon fisheries. *FAO Studies and Reviews*,
729 *GFCM* 61. FAO, Rome.
- 730 Libralato, S., Pranovi, F., Raicevich, S., Da Ponte, F., Giovanardi, O., Pastres, R., Torricelli, P., Mainardi, D.,
731 2004. Ecological stages of the Venice Lagoon analysed using landing time series data. *J.Mar.Syst.* 51,
732 331–344.
- 733 Losso, C., Volpi Ghirardini, A., 2010. Overview of ecotoxicological studies performed in the Venice Lagoon
734 (Italy). *Environ. Int.* doi:10.1016/j.envint.2009.07.017
- 735 Malavasi, S., Fiorin, R., Franco, A., Franzoi, P., Granzotto, A., Riccato, F., Mainardi, D., 2004a. Fish
736 assemblages of Venice Lagoon shallow waters: an analysis based on species, families and functional
737 guilds. *J.Mar.Syst.* 51, 19–31.
- 738 Malavasi, S., Fiorin, R., Franco, A., Torricelli, P., 2004b. Somatic energy storage and reproductive investment
739 in the grass goby *Zosterisessor ophiocephalus*. *J.Mar.Biol.Ass.U.K.* 84, 455–459.
- 740 Marchand, J., Codling, I., Drake, P., Elliott, M., Pihl, L., Rebelo, J., 2002. Environmental quality of estuaries.
741 *Fishes in Estuaries* 322–409. doi:10.1002/9780470995228.ch7
- 742 Matozzo, V., Boscolo, A., Marin, M.G., 2013. Seasonal and gender-related differences in morphometric
743 features and cellular and biochemical parameters of *Carcinus aestuarii* from the Lagoon of Venice. *Mar.*
744 *Environ. Res.* 89, 21–8. doi:10.1016/j.marenvres.2013.04.007
- 745 Matthiessen, P., Law, R.J., 2002. Contaminants and their effects on estuarine and coastal organisms in the
746 United Kingdom in the late twentieth century. *Environ. Pollut.* 120, 739–757. doi:10.1016/S0269-
747 7491(02)00175-6
- 748 McCullagh, P., Nelder, J.A., 1989. *Generalized linear models*. Second edition. London, UK.
- 749 McHugh, J.L., 1967. Estuarine nekton. *Estuaries* (Lauff, G.H., ed.). *Amer.Assoc.Adv.Sci Spec Publ.*,

- 750 Washington, DC 0, 581–619.
- 751 Molinaroli, E., Guerzoni, S., Sarretta, A., Cucco, A., Umgiesser, G., 2007. Links between hydrology and
752 sedimentology in the Lagoon of Venice, Italy. *J. Mar. Syst.* 68, 303–317.
753 doi:10.1016/j.jmarsys.2006.12.003
- 754 Molinaroli, E., Guerzoni, S., Sarretta, A., Masiol, M., Pistolato, M., 2009. Thirty-year changes (1970 to 2000)
755 in bathymetry and sediment texture recorded in the Lagoon of Venice sub-basins, Italy. *Mar. Geol.* 258,
756 115–125. doi:10.1016/j.margeo.2008.12.001
- 757 Mouillot, D., Graham, N.J., Villéger, S., Mason, N., Bellwood, D.R., 2013. A functional approach reveals
758 community responses to disturbances. *Trends Ecol. Evol.* 28, 167–77.
- 759 Mouillot, D., Laune, J., Tomasini, J.-A., Aliaume, C., Brehmer, P., Dutrieux, E., Do Chi, T., 2005. Assessment
760 of Coastal Lagoon Quality with Taxonomic Diversity Indices of Fish, Zoobenthos and Macrophyte
761 Communities. *Hydrobiologia* 550, 121–130. doi:10.1007/s10750-005-4368-y
- 762 Muxika, I., Borja, A., Bald, J., 2007. Using historical data, expert judgement and multivariate analysis in
763 assessing reference conditions and benthic ecological status, according to the European Water
764 Framework Directive. *Mar. Pollut. Bull.* 55, 16–29. doi:10.1016/j.marpolbul.2006.05.025
- 765 Nacci, D., Coiro, L., Champlin, D., Jayaraman, S., McKinney, R., Gleason, T.R., Munns, W.R., Specker, J.L.,
766 Cooper, K.R., 1999. Adaptations of wild populations of the estuarine fish *Fundulus heteroclitus* to
767 persistent environmental contaminants. *Mar. Biol.* 134, 9–17. doi:10.1007/s002270050520
- 768 Nixon, S.W., 1982. Nutrient dynamics, primary production and fisheries yields of lagoons. *Oceanol. Acta*
769 *Proceeding*, 357–371.
- 770 Noges, P., van de Bund, W., Cardoso, A.C., Solimini, A., Heiskanen, A.-S., 2009. Assessment of the ecological
771 status of European surface waters, *Developments in Hydrobiology*. Springer.
- 772 Palomares, M.L.D., Pauly, D., 2014. SeaLifeBase. World Wide Web Electron. Publ. www.sealifebase.org,
773 version (11/2014).
- 774 Pasquaud, S., Courrat, A., Fonseca, V.F., Gamito, R., Gonçalves, C.I., Lobry, J., Lepage, M., Costa, M.J.,
775 Cabral, H., 2013. Strength and time lag of relationships between human pressures and fish-based metrics
776 used to assess ecological quality of estuarine systems. *Estuar. Coast. Shelf Sci.*
777 doi:10.1016/j.ecss.2013.02.002
- 778 Pasquaud, S., Pillet, M., David, V., Sautour, B., Elie, P., 2010. Determination of fish trophic levels in an
779 estuarine system. *Estuar. Coast. Shelf Sci.* 86, 237–246.
- 780 Pérez-Domínguez, R., Maci, S., Courrat, A., Lepage, M., Borja, A., Uriarte, A., Neto, J.M., Cabral, H.,
781 St.Raykov, V., Franco, A., Alvarez, M.C., Elliott, M., 2012. Current developments on fish-based indices
782 to assess ecological-quality status of estuaries and lagoons. *Ecol. Indic.* 23, 34–45.
783 doi:10.1016/j.ecolind.2012.03.006
- 784 Pérez-Ruzafa, A., Marcos, C., 2012. Fisheries in coastal lagoons: An assumed but poorly researched aspect of
785 the ecology and functioning of coastal lagoons. *Estuar. Coast. Shelf Sci.* 110, 15–31.
786 doi:10.1016/j.ecss.2012.05.025
- 787 Pinczon Du Sel, G., Blanc, A., Daguzan, J., 2000. The diet of the cuttlefish *Sepia officinalis* L. (mollusca:
788 cephalopoda) during its life cycle in the Northern Bay of Biscay (France). *Aquat.sci.* 61, 167–178.
- 789 Potter, I.C., Tweedley, J.R., Elliott, M., Whitfield, A.K., 2013. The ways in which fish use estuaries: A

- 790 refinement and expansion of the guild approach. *Fish Fish.* 16, 1–10. doi:10.1111/faf.12050
- 791 Pranovi, F., Caccin, A., Franzoi, P., Malavasi, S., Zucchetta, M., Torricelli, P., 2013. Vulnerability of artisanal
792 fisheries to climate change in the Venice Lagoon. *J. Fish Biol.* 83, 847–864. doi:10.1111/jfb.12124
- 793 Provincia di Venezia, 2009. Piano per la gestione delle risorse alieutiche delle lagune di Venezia e Caorle. Arti
794 Grafiche Zotelli, Dosson di Casier, TV.
- 795 Provincia di Venezia, 2015. Piano per la gestione delle risorse alieutiche delle lagune di Venezia e Caorle.
796 280p. Available at: <http://pesca.provincia.venezia.it/ecm/faces/public/pesca/pprof>.
- 797 Rice, J., 2003. Environmental health indicators. *Ocean Coast. Manag.* 46, 235–259. doi:10.1016/S0964-
798 5691(03)00006-1
- 799 Rismondo, A., Curiel, D., Scarton, F., Mion, D., Caniglia, G., 2003. A new seagrass map for the Venice
800 Lagoon. Proc. 6th Int. Conf. Mediterr. Coast. Environ. MEDCOAST 03, E.Ozhan (ed.), 7-11 Oct. 2003,
801 Ravenna, Italy 0, 844–852.
- 802 Rodríguez-Climent, S., Alcaraz, C., Caiola, N., Ibáñez, C., Nebra, A., Muñoz-Camarillo, G., Casals, F.,
803 Vinyoles, D., de Sostoa, A., 2012. Gillnet selectivity in the Ebro Delta coastal lagoons and its implication
804 for the fishery management of the sand smelt, *Atherina boyeri* (Actinopterygii: Atherinidae). *Estuar.
805 Coast. Shelf Sci.* 114, 41–49. doi:10.1016/j.ecss.2011.09.008
- 806 Rozas, L.P., Minello, T.J., 1997. Estimating densities of small fishes and decapod crustaceans in shallow
807 estuarine habitats: a review of sampling design with focus on gear selection. *Estuaries* 20 (1), 199–213.
- 808 Sarretta, A., Pillon, S., Molinaroli, E., Guerzoni, S., Fontolan, G., 2010. Sediment budget in the Lagoon of
809 Venice, Italy. *Cont. Shelf Res.* 30, 934–949. doi:10.1016/j.csr.2009.07.002
- 810 Schoolmaster, D.R., Grace, J.B., Schweiger, E.W., Guntenspergen, G.R., Mitchell, B.R., Miller, K.M., Little,
811 A.M., 2013a. An algorithmic and information-theoretic approach to multimetric index construction. *Ecol.
812 Indic.* 26, 14–23. doi:10.1016/j.ecolind.2012.10.016
- 813 Schoolmaster, D.R., Grace, J.B., Schweiger, E.W., Mitchell, B.R., Guntenspergen, G.R., 2013b. A causal
814 examination of the effects of confounding factors on multimetric indices. *Ecol. Indic.* 29, 411–419.
815 doi:10.1016/j.ecolind.2013.01.015
- 816 Schoolmaster, D.R., Grace, J.B., William Schweiger, E., 2012. A general theory of multimetric indices and
817 their properties. *Methods Ecol. Evol.* 3, 773–781. doi:10.1111/j.2041-210X.2012.00200.x
- 818 Secco, T., Pellizzato, F., Sfriso, A., Pavoni, B., 2005. The changing state of contamination in the Lagoon of
819 Venice. Part 1: organic pollutants. *Chemosphere* 58, 279–90. doi:10.1016/j.chemosphere.2004.06.030
- 820 Solidoro, C., Bandelj, V., Bernardi, F., Camatti, E., Ciavatta, S., Cossarini, G., Facca, C., Franzoi, P., Libralato,
821 S., Canu, D., Pastres, R., Pranovi, F., Raicevich, S., Socal, G., Sfriso, A., Sigovini, M., Tagliapietra, D.,
822 Torricelli, P., 2010. Response of the Venice Lagoon Ecosystem to Natural and Anthropogenic Pressures
823 over the Last 50 Years, in: *Coastal Lagoons Critical Habitats of Environmental Change*. CRC Press, pp.
824 483–511. doi:10.1201/EBK1420088304-c19
- 825 Solidoro, C., Bandelj, V., Cossarini, G., Libralato, S., Melaku Canu, D., 2009. Challenges for ecological
826 modelling in a changing world: Global Changes, Sustainability and Ecosystem Based Management. *Ecol.
827 Modell.* 220, 2825–2827. doi:10.1016/j.ecolmodel.2009.08.018
- 828 Solidoro, C., Melaku canu, D., Cucco, A., Umgiesser, G., 2004. A partition of the Venice Lagoon based on
829 physical properties and analysis of general circulation. *J.Mar.Syst.* 51, 147–160.

- 830 Teatini, P., Tosi, L., Strozzi, T., Carbognin, L., Cecconi, G., Rosselli, R., Libardo, S., 2012. Resolving land
831 subsidence within the Venice Lagoon by persistent scatterer SAR interferometry. *Phys. Chem. Earth* 40-
832 41, 72–79. doi:10.1016/j.pce.2010.01.002
- 833 Umgiesser, G., Melaku canu, D., Cucco, A., Solidoro, C., 2004. A finite element model for the Venice Lagoon.
834 Development, set up, calibration and validation. *J.Mar.Syst.* 51, 123–145.
- 835 Uriarte, A., Borja, A., 2009. Assessing fish quality status in transitional waters, within the European Water
836 Framework Directive: Setting boundary classes and responding to anthropogenic pressures. *Estuar.
837 Coast. Shelf Sci.* 82, 214–224. doi:10.1016/j.ecss.2009.01.008
- 838 Vallès, H., Oxenford, H. a, 2014. The utility of simple fish community metrics for evaluating the relative
839 influence of fishing vs. other environmental drivers on Caribbean reef fish communities. *Fish Fish.* n/a-
840 n/a. doi:10.1111/faf.12085
- 841 Vasconcelos, R.P., Reis-Santos, P., Fonseca, V., Maia, A., Ruano, M., Franca, S., Vinagre, C., Costa, M.J.,
842 Cabral, H., 2007. Assessing anthropogenic pressures on estuarine fish nurseries along the Portuguese
843 coast: A multi-metric index and conceptual approach. *Sci. Total Environ.* 374, 199–215.
- 844 Wang, Y., Naumann, U., Wright, S.T., Warton, D.I., 2012. mvabund - an R package for model-based analysis
845 of multivariate abundance data. *Methods Ecol. Evol.* 3, 471–474. doi:10.1111/j.2041-
846 210X.2012.00190.x
- 847 Warton, D.I., 2008. Raw data graphing: an informative but under-utilized tool for the analysis of multivariate
848 abundances. *Austral Ecol.* 33, 290–300. doi:10.1111/j.1442-9993.2007.01816.x
- 849 Warton, D.I., Foster, S.D., De'ath, G., Stoklosa, J., Dunstan, P.K., 2014. Model-based thinking for community
850 ecology. *Plant Ecol.* doi:10.1007/s11258-014-0366-3
- 851 Warton, D.I., Wright, S.T., Wang, Y., 2012. Distance-based multivariate analyses confound location and
852 dispersion effects. *Methods Ecol. Evol.* 3, 89–101. doi:10.1111/j.2041-210X.2011.00127.x
- 853 Whitfield, A.K., Elliott, M., 2002. Fishes as indicators of environmental and ecological changes within
854 estuaries: a review of progress and some suggestions for the future. *J. Fish Biol.* 61, 229–250.
855 doi:10.1006/jfbi.2002.2079
- 856 Magistrato alle Acque di Venezia (ora Provveditorato Interregionale alle OO. PP. del Veneto- Trentino Alto
857 Adige – Friuli Venezia Giulia). 2002. Carta Tecnica della laguna di Venezia. Prodotto dal
858 Concessionario, Consorzio Venezia Nuova
- 859 Magistrato alle Acque di Venezia (ora Provveditorato Interregionale alle OO. PP. del Veneto- Trentino Alto
860 Adige – Friuli Venezia Giulia) - COSES. 2002. Campagne di rilevazione traffico lagunare 2001/2002.
861 74 p.
- 862 Magistrato alle Acque di Venezia (ora Provveditorato Interregionale alle OO. PP. del Veneto- Trentino Alto
863 Adige – Friuli Venezia Giulia) - Technital. 2002. Studio C.2.4 - Studio degli effetti della navigazione
864 interna sulla morfologia lagunare. Rapporto Finale - modello di traffico. Prodotto dal Concessionario,
865 Consorzio Venezia Nuova.
- 866 Magistrato alle Acque di Venezia (ora Provveditorato Interregionale alle OO. PP. del Veneto- Trentino Alto
867 Adige – Friuli Venezia Giulia) - Agriteco. 2004. Studio C.4.3 – III Fase: “Monitoraggio delle attività
868 alieutiche e dell’avifauna in laguna aperta”. Prodotto dal Concessionario, Consorzio Venezia Nuova.
- 869 Magistrato alle Acque di Venezia (ora Provveditorato Interregionale alle OO. PP. del Veneto- Trentino Alto

870 Adige – Friuli Venezia Giulia) - Thetis. 2004. Attività di monitoraggio ambientale della laguna di
871 Venezia MELa 1. Primo stralcio triennale (2000-2003). Prodotto dal Concessionario, Consorzio Venezia
872 Nuova.

873 Magistrato alle Acque di Venezia (ora Provveditorato Interregionale alle OO. PP. del Veneto- Trentino Alto
874 Adige – Friuli Venezia Giulia) - Thetis. 2005. Programma generale delle attività di approfondimento del
875 quadro conoscitivo di riferimento per gli interventi ambientali. 2° stralcio triennale (2003–2006)
876 “Progetto ICSEL”. Attività A. Prodotto dal Concessionario, Consorzio Venezia Nuova.

877 Magistrato alle Acque di Venezia (ora Provveditorato Interregionale alle OO. PP. del Veneto- Trentino Alto
878 Adige – Friuli Venezia Giulia) - SELC. 2005. Studio B.12.3/III. La funzionalità dell’ambiente lagunare
879 attraverso rilievi delle risorse alieutiche, dell’avifauna e dell’ittiofauna. Erodibilità del fondale e fattori di
880 disturbo:Rilievi dell'erodibilità del fondale. Rapporto intermedio. Prodotto dal Concessionario,
881 Consorzio Venezia Nuova.

882 Magistrato alle Acque di Venezia (ora Provveditorato Interregionale alle OO. PP. del Veneto- Trentino Alto
883 Adige – Friuli Venezia Giulia) - Thetis. 2005. Attività Attività di monitoraggio ambientale della laguna
884 di Venezia MELa 3. Prosecuzione delle attività di monitoraggio della qualità delle acque lagunari e loro
885 integrazione con le attività di monitoraggio e controllo condotte dalla Sezione Antinquinamento del
886 Magistrato alle Acque. Prodotto dal Concessionario, Consorzio Venezia Nuova.

887 G.R.A.L., 2006. Piano d’uso sostenibile delle aree in concessione per venericoltura. Venice (In Italian).

888 Magistrato alle Acque di Venezia (ora Provveditorato Interregionale alle OO. PP. del Veneto- Trentino Alto
889 Adige – Friuli Venezia Giulia) - Thetis. 2006. Attività di monitoraggio ambientale della laguna di
890 Venezia MELa 4. Monitoraggio di mantenimento delle conoscenze sullo stato delle acque e del
891 macrobenthos. Prodotto dal Concessionario, Consorzio Venezia Nuova.

892 Magistrato alle Acque di Venezia (ora Provveditorato Interregionale alle OO. PP. del Veneto- Trentino Alto
893 Adige – Friuli Venezia Giulia) - Agriteco. 2007. Studio B.12.3/III: “La funzionalità dell’ambiente
894 lagunare attraverso rilievi delle risorse alieutiche, dell’avifauna e dell’ittiofauna – Pesca tradizionale’.
895 Prodotto dal Concessionario, Consorzio Venezia Nuova.

896 Water Framework Directive - United Kingdom Technical Advisory Group (WFD-UKTAG, 2009. UKTAG
897 Transitional Water Assessment Methods. Fish Fauna. Transitional Fish Classification Index (TFCI).
898 SNIFFER, Scotland. www.wfduk.org ISBN: 978-1-906934-10-1

899 G.R.A.L., 2009. Adeguamento del piano d’uso sostenibile delle aree in concessione per venericoltura. Venice
900 (In Italian).

901 Magistrato alle Acque di Venezia (ora Provveditorato Interregionale alle OO. PP. del Veneto- Trentino Alto
902 Adige – Friuli Venezia Giulia) - SAMA. 2013. Dati rete di monitoraggio SAMANET.

903 www.portolando.eu

904 www.pagineazzurre.com

905 www.port-venice.com

906 www.portodichioggia.it

907 www.gralvenezia.it

908 www.fishbase.org

8. Tables

Tab. 1 Anthropogenic pressures indices and references

Category	Description	Label	References
Morphological	intertidal area lost (% area between -0.5 and +0.5 m lost between 1972 and 2002)	Intert	Sarretta et al., 2010
Morphological	gross change in bathymetry (relative change in bottom depth between 1972 and 2002, taking a variation of ± 0.75 m as minimum significant change)	Bathy	Sarretta et al., 2010
Morphological	seagrass habitat loss (relative loss between 1990 and 2002)	Seag	Caniglia et al., 1992; Rismondo et al., 2003
Morphological	relative sea level	Rise	Carbognin et al., 2010; Teatini et al., 2012
Morphological	interference with hydrographic regime (changes in sediment resuspension related to alteration of hydraulic circulation due to the presence of human infrastructures)	Hydro	D'Alpaos, 2010
resource/habitat use	aquaculture	Aqua	G.R.A.L., 2006, 2009; Provincia di Venezia, 2009
resource/habitat use	fisheries (density of fyke nets as no. traps/km ²)	Fish_class	Mag. Acque - Agriteco, 2004, 2007; Provincia di Venezia, 2009; "www.gralvenezia.it," 2015
resource/habitat use	intensity of marina developments (no. berths/km ²)	Marina	"www.pagineazzurre.com," 2014, "www.portolando.eu," 2014
resource/habitat use	navigation (no. boat passages/day)	Traffic	Mag. Acque - COSES, 2002; Mag. Acque - Technital, 2002
resource/habitat use	intensity of shipyards	Shipyard	"www.portodichioggia.it," 2014; www.port-venice.com, 2014
environmental quality	water chemical quality (the number of substances whose concentrations did not comply with imperative concentration values set for the Venice lagoon - D.M. 23/04 1998)	WatChemQual	Mag. Acque - Thetis, 2004, 2005b; ARPAV-ISPRA, 2013; Mag. Acque - SAMA, 2013
environmental quality	sediment chemical quality	SedChemQual	(Apitz et al., 2007)
environmental quality	sediment quality biological effects (Weighted Average Toxicity Index (WATI), which integrates the results of different ecotoxicological tests on different matrices)	Wati	Losso and Volpi Ghirardini, 2010
environmental quality	Benthos (Multivariate-Azti Marine Biotic Index; M-AMBI)	Mambi	Mag. Acque - Thetis, 2006; Muxika et al., 2007; ARPAV-ISPRA, 2013
environmental quality	nutrients (DIN;RP mean)	DIN; RP	Mag. Acque - Thetis, 2004, 2005b; ARPAV-ISPRA, 2013
environmental quality	Chlorophyll	Chla	Mag. Acque - Thetis, 2004, 2005b; ARPAV-ISPRA, 2013; Mag. Acque - SAMA, 2013
environmental quality	Dissolved Oxygen	Od	Mag. Acque - Thetis, 2004, 2005b; ARPAV-ISPRA, 2013; Mag. Acque - SAMA, 2013

913 **Tab. 2 Structure of models used to assess the relation between nekton community temporal factor, environmental conditions**
 914 **and anthropogenic pressures and scheme of comparison (Y_i : response variables; PC1, PC2, PC3: Principal Components of the**
 915 **PCA on environmental data; P_morpho: Pressures acting on morphology; P_Use: Pressures related to the use of the resources**
 916 **and habitat; P_Quality: Pressures affecting environmental quality of matrices; P_Fish: Pressures related to the activity of**
 917 **artisanal fishery).**

Label	Model structure /comparison	Description - Response variable affected by:
m0	$Y_i \sim \text{constant} + \epsilon_i$	None of the considered factors
m1	$Y_i \sim \text{Season X Year} + \text{constant} + \epsilon_i$	Temporal factor only
m2	$Y_i \sim \text{Season X Year} + \text{PC1} + \text{PC2} + \text{PC3} + \text{constant} + \epsilon_i$	Temporal factor and environmental conditions
m3.0	$Y_i \sim \text{Season X Year} + \text{P_Morpho} + \text{P_Use} + \text{P_Quality} + \text{constant} + \epsilon_i$	Temporal factor and anthropogenic pressures (artisanal fishery excluded)
m3.1	$Y_i \sim \text{Season X Year} + \text{PC1} + \text{PC2} + \text{PC3} + \text{P_Morpho} + \text{P_Use} + \text{P_Quality} + \text{constant} + \epsilon_i$	Temporal factor, environmental conditions and anthropogenic pressures (artisanal fishery excluded)
m4.0	$Y_i \sim \text{Season X Year} + \text{PC1} + \text{PC2} + \text{PC3} + \text{P_Fish} + \text{constant} + \epsilon_i$	Temporal factor, environmental conditions and artisanal fishery pressure
m4.1	$Y_i \sim \text{Season X Year} + \text{P_Fish} + \text{constant} + \epsilon_i$	Temporal factor and artisanal fishery pressure
m4.2	$Y_i \sim \text{Season X Year} + \text{PC1} + \text{PC2} + \text{PC3} + \text{P_Morpho} + \text{P_Use} + \text{P_Quality} + \text{P_Fish} + \text{constant} + \epsilon_i$	Temporal factor, environmental conditions, anthropogenic pressures including the one related to artisanal fishery
Testing the effect of:		
Test0	m0 vs m1	Temporal factor
Test1	m1 vs m2	Inclusion of environmental conditions if only temporal factor was considered before
Test2.0	m1 vs m3.0	Inclusion of anthropogenic pressures (artisanal fishery excluded) if only temporal factor was considered before
Test2.1	m2 vs m3.1	Inclusion of anthropogenic pressures (artisanal fishery excluded) if temporal factor and environmental conditions were considered before
Test3.0	m2 vs m4.0	Inclusion of artisanal fishery pressure if temporal factor and environmental conditions were considered before
Test3.1	m1 vs m4.1	Inclusion of artisanal fishery pressure if temporal factor was considered before
Test3.2	m3.1 vs m4.2	Inclusion of artisanal fishery pressure if temporal factor, environmental conditions and other anthropogenic pressures were considered before

918

919


920

921
922
923

Tab. 3 Summary of the significance results for the multivariate comparisons of nested models. Significant global p-values are in bold and underlined. A complete framework of the results for the multivariate comparisons is presented in supplementary materials (Table A.3), including species showing significant effect of anthropogenic pressures (if applicable).

Response variable	Test	Tested effect	p-value
Presence/Absence of all species	Test 0	Temporal factors	<u>0.001</u>
	Test 1	Environmental conditions (temporal factors already included)	p>0.05
	Test 2.0	Anthropogenic pressures (fishery excluded; temporal factors already included)	p>0.05
	Test 2.1	Anthropogenic pressures (fishery excluded; temporal factors and environmental conditions already included)	p>0.05
	Test 3.0	Fishery related pressures (temporal factors and environmental conditions already included)	p>0.05
	Test 3.1	Fishery (temporal factors already included)	p>0.05
	Test 3.2	Fishery related pressures (temporal factors, environmental conditions and other pressures already included)	p>0.05
	Biomass (Biomass of species whose cumulative biomass represent the 95% of the total)	Test 0	Temporal factors
Test 1		Environmental conditions (temporal factors already included)	<u>0.001</u>
Test 2.0		Anthropogenic pressures (fishery excluded; temporal factors already included)	<u>0.001</u>
Test 2.1		Anthropogenic pressures (fishery excluded; temporal factors and environmental conditions already included)	<u>0.001</u>
Test 3.0		Fishery related pressures (temporal factors and environmental conditions already included)	<u>0.008</u>
Test 3.1		Fishery (temporal factors already included)	p>0.05
Test 3.2		Fishery related pressures (temporal factors, environmental conditions and other pressures already included)	p>0.05


924



926

927 **Fig. 1** The Venice lagoon and the five study areas (CH: “Chioggia”; CZ: “Ca’ Zane”; LD: “Lido”; LT: “Lago
928 dei Tenei”; PL: “Ponte della Libertà”).


929



930


931 **Fig. 2** Biplots of the Principal Component Analysis on environmental variables: first two axes (A) and first
 932 and third axis (B). Biplot of the PCA on indicators of pressure: first two axes (C) and first and third axis (D).

933




934

935 **Fig. 3** Panel plot summarising the results of the nested comparison of models for the community metrics
 936 (univariate comparison): the size of the square is inversely proportional to the p-value and the shade of grey is
 937 proportional to the explained deviance. Biomass (B; left panel) and number of species (S; right panel) for
 938 fishery-related metrics (discarded, incidental, target and total catches), Estuarine Usage Functional Groups (A:
 939 anadromous; C: catadromous; ES: estuarine; ESs: solely estuarine; FS: freshwater stragglers; ME-D: marine
 940 estuarine-dependent; ME-O: marine estuarine-opportunists; MS: marine stragglers) and Feeding Mode
 941 Functional Groups (Bma: macrobenthivores; Bmi: microbenthivores; DV: detritivores; HP:
 942 hyperbenthivores/piscivores; HV: herbivores; HZ: hyperbenthivores/zooplanktivores; OV: omnivores; PL:
 943 planktivores). For each comparison test, the effects tested are indicated as follows: t = temporal factor; e =
 944 environmental conditions; p = anthropogenic pressures (other than fishery); f = artisanal fishery pressure.



945

946 **Fig. 4** Averaged coefficients (\pm C. I.) for the 4 considered pressure categories (x-axis) for the models of biomass
 947 (B; left column of plots) and number of species (S; right panel of plots) for (a) fishery-related metrics
 948 (discarded, incidental, target and total catches), (b) Estuarine Usage Functional Groups (A: anadromous; C:
 949 catadromous; ES: estuarine; ESs: solely estuarine; FS: freshwater stragglers; ME-D: marine estuarine-
 950 dependent; ME-O: marine estuarine-opportunists; MS: marine stragglers) and (c) Feeding Mode Functional
 951 Groups (Bma: macrobenthivores; Bmi: microbenthivores; DV: detritivores; HP: hyperbenthivores/piscivores;
 952 HV: herbivores; HZ: hyperbenthivores/zooplanktivores; OV: omnivores; PL: planktivores). The size of the
 953 dots are proportional to the Akaike weights used for model averaging.



954

955 **Fig. 5** Averaged coefficients (\pm S. E.) for the four considered pressure categories (x-axis) for the models of
 956 presence/absence (left panel) and biomass (right panel) for all considered species.